

09230817

Trying 3106016892...Open

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1600LXD

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
NEWS 2 Dec 17 The CA Lexicon available in the CAPLUS and CA files
NEWS 3 Feb 06 Engineering Information Encompass files have new names
NEWS 4 Feb 16 TOXLINE no longer being updated
NEWS 5 Apr 23 Search Derwent WPINDEX by chemical structure
NEWS 6 Apr 23 PRE-1967 REFERENCES NOW SEARCHABLE IN CAPLUS AND CA
NEWS 7 May 07 DGENE Reload
NEWS 8 Jun 20 Published patent applications (A1) are now in USPATFULL
NEWS 9 JUL 13 New SDI alert frequency now available in Derwent's
DWPI and DPCI
NEWS 10 Aug 23 In-process records and more frequent updates now in
MEDLINE
NEWS 11 Aug 23 PAGE IMAGES FOR 1947-1966 RECORDS IN CAPLUS AND CA
NEWS 12 Aug 23 Adis Newsletters (ADISNEWS) now available on STN
NEWS 13 Sep 17 IMSworld Pharmaceutical Company Directory name change
to PHARMASEARCH
NEWS 14 Oct 09 Korean abstracts now included in Derwent World Patents
Index
NEWS 15 Oct 09 Number of Derwent World Patents Index updates increased
NEWS 16 Oct 15 Calculated properties now in the REGISTRY/ZREGISTRY File
NEWS 17 Oct 22 Over 1 million reactions added to CASREACT
NEWS 18 Oct 22 DGENE GETSIM has been improved

NEWS EXPRESS August 15 CURRENT WINDOWS VERSION IS V6.0c,
CURRENT MACINTOSH VERSION IS V6.0 (ENG) AND V6.0J (JP),
AND CURRENT DISCOVER FILE IS DATED 07 AUGUST 2001
NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS INTER General Internet Information
NEWS LOGIN Welcome Banner and News Items
NEWS PHONE Direct Dial and Telecommunication Network Access to STN
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that
specific topic.

All use of STN is subject to the provisions of the STN Customer
agreement. Please note that this agreement limits use to scientific
research. Use for software development or design or implementation
of commercial gateways or other similar uses is prohibited and may
result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 06:19:58 ON 24 OCT 2001

09230817

=> file reg

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.15

0.15

FILE 'REGISTRY' ENTERED AT 06:20:13 ON 24 OCT 2001

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2001 American Chemical Society (ACS)

STRUCTURE FILE UPDATES: 22 OCT 2001 HIGHEST RN 364037-88-7

DICTIONARY FILE UPDATES: 22 OCT 2001 HIGHEST RN 364037-88-7

TSCA INFORMATION NOW CURRENT THROUGH July 7, 2001

Please note that search-term pricing does apply when
conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER see
HELP CROSSOVER for details.

Calculated physical property data is now available. See HELP PROPERTIES
for more information. See STNote 27, Searching Properties in the CAS
Registry File, for complete details:

<http://www.cas.org/ONLINE/STN/STNOTES/stnotes27.pdf>

=> s valsartan/CN

L1 1 VALSARTAN/CN

=> s L1

L2 1 VALSARTAN/CN

=> file caplus

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

8.22

8.37

FILE 'CAPLUS' ENTERED AT 06:21:13 ON 24 OCT 2001

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2001 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is
held by the publishers listed in the PUBLISHER (PB) field (available
for records published or updated in Chemical Abstracts after December
26, 1996), unless otherwise indicated in the original publications.

FILE COVERS 1947 - 24 Oct 2001 VOL 135 ISS 18

FILE LAST UPDATED: 22 Oct 2001 (20011022/ED)

This file contains CAS Registry Numbers for easy and accurate
substance identification.

This file supports REGlstry for direct browsing and searching of
all substance data from the REGISTRY file. Enter HELP FIRST for
more information.

09230817

CAPLUS now provides online access to patents and literature covered in CA from 1947 to the present. On April 22, 2001, bibliographic information and abstracts were added for over 2.2 million references published in CA from 1947 to 1966.

The CA Lexicon is now available in the Controlled Term (/CT) field. Enter HELP LEXICON for full details.

Attention, the CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

=> s l1

L3 233 L1

=> s l3 and structure

2009554 STRUCTURE

530668 STRUCTURES

2258454 STRUCTURE

(STRUCTURE OR STRUCTURES)

L4 9 L3 AND STRUCTURE

=> d l4 1-9 ibib abs abs

L4 ANSWER 1 OF 9 CAPLUS COPYRIGHT 2001 ACS

ACCESSION NUMBER: 2001:311121 CAPLUS

DOCUMENT NUMBER: 135:251404

TITLE: Does protein binding modulate the effect of angiotensin II receptor antagonists?

AUTHOR(S): Maillard, Marc P.; Centeno, Catherine; Frostell-Karlsson, Asa; Brunner, Hans R.; Burnier, Michel

CORPORATE SOURCE: Division of Hypertension and Vascular Medicine, Lausanne University Hospital, Lausanne, CH-1011, Switz.

SOURCE: JRAAS (2001), 2(Suppl. 1), S54-S58

CODEN: JRAAAG; ISSN: 1470-3203

PUBLISHER: JRAAS Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Angiotensin II AT1-receptor antagonists are highly bound to blood plasma proteins (.gtoreq. 99%). With some antagonists, such as DuP-532, the protein binding was such that no efficacy of the drug could be demonstrated clin. Whether protein binding interferes with the efficacy of other antagonists is not known. The authors have therefore investigated in vitro how plasma proteins may affect the antagonistic effect of different AT1-receptor antagonists. A radio-receptor binding assay was used to analyze the interaction between proteins and the

ability

of various angiotensin II (Ang II) antagonists to block AT1-receptors.

In

addn., the Biacore technol., a new technique which enables the real-time monitoring of binding events between 2 mols., was used to evaluate the dissocn. rate consts. of 5 AT1-receptor antagonists from human serum albumin. The in vitro AT1-antagonistic effects of different Ang II receptor antagonists were differentially affected by the presence of

human

plasma, with rightward shifts of the IC50 ranging from 1 to several orders of magnitude. The importance of the shift correlates with the dissocn. rate consts. of these drugs from albumin. The authors' expts. also show that the way that AT1-receptor antagonists bind to proteins differs from

1 compd. to another. These results suggest that the interaction with plasma

proteins appears to modulate the efficacy of some Ang II antagonists. Although the high binding level of Ang II receptor antagonist to plasma proteins appears to be a feature common to this class of compds., the kinetics and characteristics of this binding is of great importance.

With some antagonists, protein binding interferes markedly with their efficacy to block AT1-receptors.

AB Angiotensin II AT1-receptor antagonists are highly bound to blood plasma proteins (.gtoreq. 99%). With some antagonists, such as DuP-532, the protein binding was such that no efficacy of the drug could be demonstrated clin. Whether protein binding interferes with the efficacy of other antagonists is not known. The authors have therefore investigated in vitro how plasma proteins may affect the antagonistic effect of different AT1-receptor antagonists. A radio-receptor binding assay was used to analyze the interaction between proteins and the

ability of various angiotensin II (Ang II) antagonists to block AT1-receptors.

In addn., the Biacore technol., a new technique which enables the real-time monitoring of binding events between 2 mols., was used to evaluate the dissocn. rate consts. of 5 AT1-receptor antagonists from human serum albumin. The in vitro AT1-antagonistic effects of different Ang II receptor antagonists were differentially affected by the presence of

human plasma, with rightward shifts of the IC50 ranging from 1 to several orders of magnitude. The importance of the shift correlates with the dissocn. rate consts. of these drugs from albumin. The authors' expts. also show that the way that AT1-receptor antagonists bind to proteins differs from

1 compd. to another. These results suggest that the interaction with plasma

proteins appears to modulate the efficacy of some Ang II antagonists. Although the high binding level of Ang II receptor antagonist to plasma proteins appears to be a feature common to this class of compds., the kinetics and characteristics of this binding is of great importance.

With some antagonists, protein binding interferes markedly with their efficacy to block AT1-receptors.

REFERENCE COUNT: 12

REFERENCE(S): (1) Burnier, M; Lancet 2000, V355, P637 CAPLUS
(2) Chiu, A; Biochem Biophys Res Comm 1991, V177(1), P209 CAPLUS
(3) Csajka, C; Pharmacokinet 1997, V32, P1 CAPLUS
(4) Goldberg, M; Clin Pharmacol ther 1997, V61, P59 CAPLUS
(6) Maillard, M; Am J Hypertens 1999, V12, P1201 CAPLUS

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 2 OF 9 CAPLUS COPYRIGHT 2001 ACS

ACCESSION NUMBER: 2001:82589 CAPLUS

DOCUMENT NUMBER: 135:86860

TITLE: Antiplatelet action of losartan involves TXA2 receptor

AUTHOR(S): antagonism but not TXA2 synthase inhibition
Chlopicki, S.; Koda, M.; Chabielska, E.; Buczek, W.; Gryglewski, R. J.CORPORATE SOURCE: Department of Pharmacology Medical College,
Jagiellonian University, Pol.SOURCE: J. Physiol. Pharmacol. (2000), 51(4, Pt. 1), 715-722
CODEN: JPHPEI; ISSN: 0867-5910

PUBLISHER: Polish Physiological Society

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Various AT1 receptor antagonists including losartan are known to inhibit human platelet activation by antagonizing TXA2/PGH2 receptors (TP receptors). Presently, we check a hypothesis that losartan, an imidazole deriv. in contrast with valsartan, a non-imidazole compd., may inhibit human platelet activation also through inhibition of TXA2 synthesis. Inhibitory action of losartan, its active metabolite EXP 3174, and valsartan, on collagen-induced platelet aggregation and TXA2 generation was compared to effects achieved by each compd. on U 46619-induced aggregation in aspirinized platelets. Losartan and aspirin inhibited collagen-induced platelet aggregation with approx. the same potency, whereas EXP 3174 and valsartan showed much weaker antiplatelet effects. Interestingly, losartan, EXP 3174 and valsartan displayed similar potencies as inhibitors of U 46619-induced aggregation in aspirinized platelets as in collagen-induced aggregation in non-aspirinized platelets.

None of the above three AT1 antagonists, up to a concn. of 300 .mu.M, did influence collagen-induced TXA2 synthesis in human platelets. In conclusion, antiplatelet effects of AT1 antagonists, irresp. of the presence or absence of non-condensed imidazole in their chem.

structure, involve antagonism of TP receptors but not inhibition of TXA2 synthesis in platelets.

AB Various AT1 receptor antagonists including losartan are known to inhibit human platelet activation by antagonizing TXA2/PGH2 receptors (TP receptors). Presently, we check a hypothesis that losartan, an imidazole deriv. in contrast with valsartan, a non-imidazole compd., may inhibit human platelet activation also through inhibition of TXA2 synthesis. Inhibitory action of losartan, its active metabolite EXP 3174, and valsartan, on collagen-induced platelet aggregation and TXA2 generation was compared to effects achieved by each compd. on U 46619-induced aggregation in aspirinized platelets. Losartan and aspirin inhibited collagen-induced platelet aggregation with approx. the same potency, whereas EXP 3174 and valsartan showed much weaker antiplatelet effects. Interestingly, losartan, EXP 3174 and valsartan displayed similar potencies as inhibitors of U 46619-induced aggregation in aspirinized platelets as in collagen-induced aggregation in non-aspirinized platelets.

None of the above three AT1 antagonists, up to a concn. of 300 .mu.M, did influence collagen-induced TXA2 synthesis in human platelets. In conclusion, antiplatelet effects of AT1 antagonists, irresp. of the presence or absence of non-condensed imidazole in their chem.

structure, involve antagonism of TP receptors but not inhibition of TXA2 synthesis in platelets.

L4 ANSWER 3 OF 9 CAPLUS COPYRIGHT 2001 ACS

ACCESSION NUMBER: 2000:805989 CAPLUS

DOCUMENT NUMBER: 134:101153

TITLE: Carbon-14 labelling of DIOVAN in its valine-moiety

AUTHOR(S): Moenius, Th.; Voges, R.; Burtscher, P.; Zuger, Ch.

CORPORATE SOURCE: Preclinical Safety, DMPK-IL, Novartis Pharma AG, CH-Basel, Switz.

SOURCE: J. Labelled Compd. Radiopharm. (2000), 43(13), 1245-1252


CODEN: JLCRD4; ISSN: 0362-4803

PUBLISHER: John Wiley & Sons Ltd.

DOCUMENT TYPE: Journal


LANGUAGE: English

GI



AB As a highly specific and non peptide AT₁-antagonist Valsartan (I) is marketed under the trade-name DIOVAN for effective treatment of hypertension. This paper describes the synthesis of ¹⁴C-labeled I, which incorporates two ¹⁴C isotopes in the valine-moiety. Reaction of (-)-bromo-[1,2-¹⁴C]acetyl bornane-10.2-sultam ((-)-[¹⁴C₂]BABS) with benzophenone imine gave (-)-diphenyl-methylene[1,2-¹⁴C₂]glycinyl bornane-10.2-sultam ((-)-[¹⁴C₂]DPMGBS), which was alkylated with 2-iodopropane to build-up the valine **structure**. Initially the resulting sultam-protected valine (II) was treated with treated with bromide (III) to produce the precursor. However, under conditions routinely used for sultam-cleavage deprotection resulted in the racemization of the amino acid. Successful cleavage was accomplished via N-Boc-protection of II followed by hydrolytic cleavage of the auxiliary and esterification to give the L-[¹⁴C₂]valine benzyl ester. Finally [¹⁴C₂]-I was synthesized in a 10 step synthesis in an overall radiochem. yield of 10% relative to the (-)-[1,2-¹⁴C]BABS employed.

GI



AB As a highly specific and non peptide AT₁-antagonist Valsartan (I) is marketed under the trade-name DIOVAN for effective treatment of hypertension. This paper describes the synthesis of ¹⁴C-labeled I, which incorporates two ¹⁴C isotopes in the valine-moiety. Reaction of (-)-bromo-[1,2-¹⁴C]acetyl bornane-10.2-sultam ((-)-[¹⁴C₂]BABS) with benzophenone imine gave (-)-diphenyl-methylene[1,2-¹⁴C₂]glyciny bornane-10.2-sultam ((-)-[¹⁴C₂]DPMGBS), which was alkylated with 2-iodopropane to build-up the valine **structure**. Initially the resulting sultam-protected valine (II) was treated with treated with bromide (III) to produce the precursor. However, under conditions routinely used for sultam-cleavage deprotection resulted in the racemization of the amino acid. Successful cleavage was accomplished via N-Boc-protection of II followed by hydrolytic cleavage of the auxiliary and esterification to give the L-[¹⁴C₂]valine benzyl ester. Finally [¹⁴C₂]-I was synthesized in a 10 step synthesis in an overall radiochem. yield of 10% relative to the (-)-[1,2-¹⁴C]BABS employed.

REFERENCE COUNT: 14
 REFERENCE(S): (2) Anon; US 4355040 CAPLUS
 (4) Buhlmayer, P; Bioorganic & Medicinal Chemistry Letters 1994, V4, P29 CAPLUS
 (5) Burtscher, P; J Labelled Cpd Radiopharm 2000, V43, P205 CAPLUS
 (6) Criscione, L; Cardiovascular Drug Reviews 1995, V13, P230 CAPLUS
 (7) de Gasparo, M; Regulatory Peptides 1995, V59, P303 CAPLUS
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 4 OF 9 CAPLUS COPYRIGHT 2001 ACS
 ACCESSION NUMBER: 2000:476348 CAPLUS
 DOCUMENT NUMBER: 133:171982
 TITLE: Basilar artery remodelling in the genetically hypertensive rat: effects of nitric oxide synthase

inhibition and treatment with valsartan and enalapril
 AUTHOR(S): Ledingham, J. M.; Laverty, R.
 CORPORATE SOURCE: Department of Pharmacology, School of Medical
 Sciences, University of Otago, Dunedin, N. Z.
 SOURCE: Clin. Exp. Pharmacol. Physiol. (2000), 27(8), 642-646
 CODEN: CEXPB9; ISSN: 0305-1870
 PUBLISHER: Blackwell Science Asia Pty Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB 1. The **structure** of the basilar artery and the relationship of **structure** to blood pressure and ventricular hypertrophy was examd. in genetically hypertensive (GH) rats, their control normotensive (N) Wistar strain, GH given the nitric oxide synthase (NOS) inhibitor, NG-nitro-L-arginine Me ester (L-NAME) and GH given L-NAME and either valsartan or enalapril. 2. Systolic blood pressure (SBP; tail-cuff) was measured weekly from age 7-12 wk. At the end of the expt. at 12 wk, the basilar artery was fixed by perfusion and embedded in Technovit (Heraeus Kulzer GmbH, Werheim, Germany). Serial sections were cut and stained and stereol. anal. applied to quantify the morphol. of the vessels. Left ventricular (LV) mass was detd. 3. Both SBP and LV mass were significantly increased in GH compared with N ($P < 0.001$) and increased further in GH given L-NAME ($P < 0.05$). The GH L-NAME + valsartan and GH L-NAME + enalapril groups had significantly lower ($P < 0.001$) SBP and LV mass than the GH L-NAME group. 4. Basilar arteries in GH (which are frankly hypertensive, but have no apparent endothelial defect) showed hypotrophic inward remodelling compared with the N control group with no change in media to lumen ratio. 5. In the GH L-NAME group, further inward remodelling occurred and the media to lumen ratio was increased compared with N ($P < 0.01$) and GH ($P < 0.05$). Valsartan treatment in GH L-NAME rats caused eutrophic outward remodelling. Enalapril caused hypertrophic outward remodelling, suggesting that the angiotensin II-stimulated growth was not entirely suppressed with an angiotensin-converting enzyme inhibitor or that there was a bradykinin effect with enalapril. 6. In GH with an endothelial defect induced by treatment with L-NAME, the further remodelling, together with an increased media to lumen ratio and the development of a stroke-like syndrome, indicates the NOS-inhibited GH rat may be a useful model for essential hypertension (where it is known that endothelial abnormalities exist) and where stroke can develop as a consequence of the hypertension.

AB 1. The **structure** of the basilar artery and the relationship of **structure** to blood pressure and ventricular hypertrophy was examd. in genetically hypertensive (GH) rats, their control normotensive (N) Wistar strain, GH given the nitric oxide synthase (NOS) inhibitor, NG-nitro-L-arginine Me ester (L-NAME) and GH given L-NAME and either valsartan or enalapril. 2. Systolic blood pressure (SBP; tail-cuff) was measured weekly from age 7-12 wk. At the end of the expt. at 12 wk, the basilar artery was fixed by perfusion and embedded in Technovit (Heraeus Kulzer GmbH, Werheim, Germany). Serial sections were cut and stained and stereol. anal. applied to quantify the morphol. of the vessels. Left ventricular (LV) mass was detd. 3. Both SBP and LV mass were significantly increased in GH compared with N ($P < 0.001$) and increased further in GH given L-NAME ($P < 0.05$). The GH L-NAME + valsartan and GH L-NAME + enalapril groups had significantly lower ($P < 0.001$) SBP and LV mass than the GH L-NAME group. 4. Basilar arteries in GH (which are frankly hypertensive, but have no apparent endothelial defect) showed hypotrophic inward remodelling compared with the N control group with no change in media to lumen ratio. 5. In the GH L-NAME group, further inward

09230817

remodelling occurred and the media to lumen ratio was increased compared with N ($P < 0.01$) and GH ($P < 0.05$). Valsartan treatment in GH L-NAME rats caused eutrophic outward remodelling. Enalapril caused hypertrophic outward remodelling, suggesting that the angiotensin II-stimulated growth was not entirely suppressed with an angiotensin-converting enzyme inhibitor or that there was a bradykinin effect with enalapril. 6. In GH with an endothelial defect induced by treatment with L-NAME, the further remodelling, together with an increased media to lumen ratio and the development of a stroke-like syndrome, indicates the NOS-inhibited GH rat may be a useful model for essential hypertension (where it is known that endothelial abnormalities exist) and where stroke can develop as a consequence of the hypertension.

REFERENCE COUNT: 27
REFERENCE(S): (3) Blot, S; Stroke 1994, V25, P1666 CAPLUS
(4) Diederich, D; Ant J Physiol 1990, V258, PH445 CAPLUS
(11) Konishi, M; Hypertension 1983, V5, P881 CAPLUS
(12) Ledingham, J; Clin Exp Pharmacol Physiol 1997, V24, P433 CAPLUS
(14) Lee, R; Pharmacol Ther 1995, V66, P149 CAPLUS
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 5 OF 9 CAPLUS COPYRIGHT 2001 ACS

ACCESSION NUMBER: 1997:204117 CAPLUS
DOCUMENT NUMBER: 126:195243
TITLE: Method for treating renal disease using an ACE inhibitor and an angiotensin II antagonist
INVENTOR(S): Remuzzi, Giuseppe; Eydeloth, Ronald S.; Owen, Roger A.; Shahinfar, Shahnaz
PATENT ASSIGNEE(S): Merck and Co., Inc., USA; Laboratoires Merck Sharp & Dohme-Chibret Snc; Remuzzi, Giuseppe; Eydeloth, Ronald S.; Owen, Roger A.; Shahinfar, Shahnaz
SOURCE: PCT Int. Appl., 32 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9702032	A1	19970123	WO 1996-US10942	19960626
W:	AL, AM, AU, AZ, BB, BG, BR, BY, CA, CN, CZ, EE, GE, HU, IL, IS, JP, KG, KR, KZ, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TR, TT, UA, US, UZ, VN, AM, AZ, BY, KG, KZ			
RW:	KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG			
CA 2224451	AA	19970123	CA 1996-2224451	19960626
AU 9662916	A1	19970205	AU 1996-62916	19960626
AU 716519	B2	20000224		
EP 835106	A1	19980415	EP 1996-921794	19960626
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, PT, IE, FI			
JP 11508894	T2	19990803	JP 1996-505200	19960626
PRIORITY APPLN. INFO.:			US 1995-770	19950630
			GB 1996-2854	19960213

WO 1996-US10942 19960626

AB The present invention relates to a method of treating and/or preventing renal disease with the coadministration of an ACE inhibitor and an AII receptor antagonist. The present invention also relates to a method for protection of renal **structure** and/or renal function with the coadministration of an ACE inhibitor and an AII receptor antagonist. The combination is also useful in preventing renal injury and protecting glomerular **structure**. The effect of Lisinopril (ACE inhibitor) and Losartan (angiotensin II receptor antagonist) in animals with

diabetic
nephropathy is described.

AB The present invention relates to a method of treating and/or preventing renal disease with the coadministration of an ACE inhibitor and an AII receptor antagonist. The present invention also relates to a method for protection of renal **structure** and/or renal function with the coadministration of an ACE inhibitor and an AII receptor antagonist. The combination is also useful in preventing renal injury and protecting glomerular **structure**. The effect of Lisinopril (ACE inhibitor) and Losartan (angiotensin II receptor antagonist) in animals with

diabetic
nephropathy is described.

L4 ANSWER 6 OF 9 CAPLUS COPYRIGHT 2001 ACS

ACCESSION NUMBER: 1997:50392 CAPLUS

DOCUMENT NUMBER: 126:139974

TITLE: A 3D QSAR CoMFA study of non-peptide angiotensin II receptor antagonists

AUTHOR(S): Belvisi, Laura; Bravi, Gianpaolo; Catalano, Giovanna; Mabilia, Massimo; Salimbeni, Aldo; Scolastico, Carlo

CORPORATE SOURCE: Organic and Industrial Chemistry Department, C.N.R. (National Research Council) Centre for the Study of Organic and Natural Compounds, University of Milan, Milan, I-20133, Italy

SOURCE: J. Comput.-Aided Mol. Des. (1996), 10(6), 567-582

CODEN: JCADEQ; ISSN: 0920-654X

PUBLISHER: ESCOM

DOCUMENT TYPE: Journal

LANGUAGE: English

AB A series of non-peptide angiotensin II receptor antagonists was investigated with the aim of developing a 3D QSAR model using comparative mol. field anal. descriptors and approaches. The main goals of the study were dictated by an interest in methodologies and an understanding of the binding requirements to the AT1 receptor. Consistency with the

previously
derived activity models was always checked to contemporarily test the validity of the various hypotheses. The specific conformations chosen

for
the study, the procedures invoked to superimpose all **structures**, the conditions employed to generate steric and electrostatic field values and the various PCA/PLS runs are discussed in detail. The effect of exptl. design techniques to select objects (mols.) and variables (descriptors) with respect to the predictive power of the QSAR models derived was esp. analyzed.

AB A series of non-peptide angiotensin II receptor antagonists was investigated with the aim of developing a 3D QSAR model using comparative mol. field anal. descriptors and approaches. The main goals of the study were dictated by an interest in methodologies and an understanding of the binding requirements to the AT1 receptor. Consistency with the

previously

09230817

derived activity models was always checked to contemporarily test the validity of the various hypotheses. The specific conformations chosen for the study, the procedures invoked to superimpose all **structures**, the conditions employed to generate steric and electrostatic field values and the various PCA/PLS runs are discussed in detail. The effect of exptl. design techniques to select objects (mols.) and variables (descriptors) with respect to the predictive power of the QSAR models derived was esp. analyzed.

L4 ANSWER 7 OF 9 CAPLUS COPYRIGHT 2001 ACS

ACCESSION NUMBER: 1997:29751 CAPLUS

DOCUMENT NUMBER: 126:98857

TITLE: Non-peptide angiotensin II receptor antagonists: PCA &

PLS analyses employing different types of 3D molecular

descriptors

AUTHOR(S): Mabilia, M.; Fioravanzo, E.

CORPORATE SOURCE: S.IN - Soluzioni Informatiche, Vicenza, I-36100, Italy

SOURCE: Bioact. Compd. Des. (1996), 127-136. Editor(s): Ford,

Martyn G. Bios Scientific Publishers: Oxford, UK.

CODEN: 63SXAI

DOCUMENT TYPE: Conference

LANGUAGE: English

AB Different three-dimensional (3D) mol. descriptors were generated for a set

of angiotensin II receptor antagonists with the aim to: a. establish quant. **structure**-activity relationships (QSAR) by means of Principal Component Anal. (PCA) and Partial Least-Squares (PLS) methods, b. compare the modeling and predictive capabilities of the different

types of descriptors considered, c. establish the effect of exptl. design techniques to select objects (mols.) and variables (descriptors), with respect to the predictive power of the QSAR models derived.

AB Different three-dimensional (3D) mol. descriptors were generated for a set

of angiotensin II receptor antagonists with the aim to: a. establish quant. **structure**-activity relationships (QSAR) by means of Principal Component Anal. (PCA) and Partial Least-Squares (PLS) methods, b. compare the modeling and predictive capabilities of the different

types of descriptors considered, c. establish the effect of exptl. design techniques to select objects (mols.) and variables (descriptors), with respect to the predictive power of the QSAR models derived.

L4 ANSWER 8 OF 9 CAPLUS COPYRIGHT 2001 ACS

ACCESSION NUMBER: 1996:381984 CAPLUS

DOCUMENT NUMBER: 125:75808

TITLE: Remodelling of resistance arteries in genetically hypertensive rats by treatment with valsartan, an angiotensin II receptor antagonist

AUTHOR(S): Ledingham, Janet M.; Lavery, R.

CORPORATE SOURCE: Dep. Pharmacol., Univ. Otago Med. Sch., Dundee, N. Z.

SOURCE: Clin. Exp. Pharmacol. Physiol. (1996), 23(6/7),

DOCUMENT TYPE:

Journal

LANGUAGE:

English

AB In the present study, New Zealand genetically hypertensive (GH) rats were treated with valsartan, a specific angiotensin II (AT1) receptor antagonist, to measure the effects on blood pressure (BP), cardiac hypertrophy and the **structure** of resistance arteries. Normotensive (N) rats were used as controls. Valsartan (val) was given

to

GH rats at three different doses (10, 3 or 0.3 mg/kg per day, via osmotic mini-pumps implanted i.p.) from age 4-10 wk. Untreated GH (mini-pump + vehicle) and N rats (mini-pump + vehicle) were used as controls. BP was measured weekly and at the end of the expt., left ventricular (LV) mass was recorded and the **structure** of mesenteric resistance arteries (MRA) was detd. using stereol. methods. BP fell in a dose-dependent fashion, being reduced to normotensive levels by 10 mg/kg; LV mass was significantly reduced ($P < 0.0001$) to below normotensive values in the GH group given 10 mg/kg val and significantly reduced ($P < 0.001$), although not normalized, in the other two treatment groups. In MRA, the media/lm (M/L) ratio was reduced by val according to dose level, to below normotensive values in GHval10, and to levels not different from normotensive values in the GHval3 and GHval0.3 groups. The hypertrophy

of

smooth muscle cells in GH rats was reduced with val treatment at all doses. Reversal of cardiac and vascular hypertrophy occurred even when

BP

was not reduced to normotensive levels, indicating an effect on vessel growth but without any retardation of body growth. These results suggest that the vascular structural changes seen after val and angiotensin converting enzyme inhibitor treatment are probably due to the blocking of angiotensin rather than any bradykinin effect.

AB In the present study, New Zealand genetically hypertensive (GH) rats were treated with valsartan, a specific angiotensin II (AT1) receptor antagonist, to measure the effects on blood pressure (BP), cardiac hypertrophy and the **structure** of resistance arteries. Normotensive (N) rats were used as controls. Valsartan (val) was given

to

GH rats at three different doses (10, 3 or 0.3 mg/kg per day, via osmotic mini-pumps implanted i.p.) from age 4-10 wk. Untreated GH (mini-pump + vehicle) and N rats (mini-pump + vehicle) were used as controls. BP was measured weekly and at the end of the expt., left ventricular (LV) mass was recorded and the **structure** of mesenteric resistance arteries (MRA) was detd. using stereol. methods. BP fell in a dose-dependent fashion, being reduced to normotensive levels by 10 mg/kg; LV mass was significantly reduced ($P < 0.0001$) to below normotensive values in the GH group given 10 mg/kg val and significantly reduced ($P < 0.001$), although not normalized, in the other two treatment groups. In MRA, the media/lm (M/L) ratio was reduced by val according to dose level, to below normotensive values in GHval10, and to levels not different from normotensive values in the GHval3 and GHval0.3 groups. The hypertrophy

of

smooth muscle cells in GH rats was reduced with val treatment at all doses. Reversal of cardiac and vascular hypertrophy occurred even when

BP

was not reduced to normotensive levels, indicating an effect on vessel growth but without any retardation of body growth. These results suggest that the vascular structural changes seen after val and angiotensin

09230817

converting enzyme inhibitor treatment are probably due to the blocking of angiotensin rather than any bradykinin effect.

L4 ANSWER 9 OF 9 CAPLUS COPYRIGHT 2001 ACS

ACCESSION NUMBER: 1994:260544 CAPLUS

DOCUMENT NUMBER: 120:260544

TITLE: Valsartan, a potent, orally active angiotensin II antagonist developed from the structurally new amino acid series

AUTHOR(S): Buhlmayer, Peter; Furet, Pascal; Criscione, Leoluca; de Gasparo, Marc; Whitebread, Steven; Schmidlin, Tibur; Lattmann, Rene; Wood, Jeanette

CORPORATE SOURCE: Res. Dep., Ciba-Geigy Ltd., Basel, CH-4002, Switz.


SOURCE: Bioorg. Med. Chem. Lett. (1994), 4(1), 29-34

CODEN: BMCLE8; ISSN: 0960-894X

DOCUMENT TYPE: Journal


LANGUAGE: English

GI



AB Starting from the **structure** of DuP-753 (I) and a 3-dimensional model of the pentapeptide Tyr-Ile-His-Pro-Ile, a series of new and highly potent antagonist has been designed where the imidazole moiety of the DuPont compd. has been replaced by an N-acylated aminoacid residue. Valsartan (CGP48933) [(S)-N-valeryl-N-[[2'-(1H-tetrazol-5-yl)biphenyl-4-yl]methyl]valine], has been selected for clin. investigation.

GI



AB Starting from the **structure** of DuP-753 (I) and a 3-dimensional

09230817

model of the pentapeptide Tyr-Ile-His-Pro-Ile, a series of new and highly potent antagonist has been designed where the imidazole moiety of the DuPont compd. has been replaced by an N-acylated aminoacid residue. Valsartan (CGP48933) [(S)-N-valeryl-N-[[2'-(1H-tetrazol-5-yl)biphenyl-4-yl]methyl]valine], has been selected for clin. investigation.

=> file stnguide

COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
22.42	30.79

FULL ESTIMATED COST

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE	TOTAL
ENTRY	SESSION
-5.29	-5.29

CA SUBSCRIBER PRICE

FILE 'STNGUIDE' ENTERED AT 06:23:43 ON 24 OCT 2001
USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
COPYRIGHT (C) 2001 AMERICAN CHEMICAL SOCIETY, JAPAN SCIENCE
AND TECHNOLOGY CORPORATION, AND FACHINFORMATIONSZENTRUM KARLSRUHE

FILE CONTAINS CURRENT INFORMATION.
LAST RELOADED: Oct 19, 2001 (20011019/UP).